

xenogeneic DNA segment of at least 100 kb which method comprises:

combining under fusing conditions said
available embryonic stem cells and yeast spheroplasts, said
5 spheroplasts containing a yeast artificial chromosome (YAC)
comprising said xenogeneic DNA segment and including a marker
for selection, whereby said xenogeneic DNA segment becomes
stably integrated into the genome of said embryonic stem
cells; and

selecting for an embryonic stem cell carrying
said xenogeneic DNA segment by means of the marker.

2. (Amended) An improved method for producing a
modified nonhuman animal, said animal having a nonyeast
15 xenogeneic DNA segment of at least 100 kb stably integrated
into the genome of at least some of the germ cells of said
animal, said method comprising:

combining under fusing conditions embryonic
stem (ES) cells of an available ES cell line derived from said
20 animal and yeast spheroplasts, said spheroplasts containing a
yeast artificial chromosome (YAC) comprising said xenogeneic
DNA segment and including a marker for selection, whereby said
xenogeneic DNA segment becomes integrated into the genome of
said embryonic stem cells;

25 selecting for embryonic stem cells carrying
said xenogeneic DNA segment by means of the marker; and

transferring said embryonic cells into a host
blastocyst, implanting said blastocyst in a pseudopregnant
animal recipient, and allowing said blastocyst to develop to
30 term to produce a chimeric animal carrying said xenogeneic DNA
segment integrated into the genome of at least some of the
germ cells of said animal.

3. (Amended) An improved method for producing a modified nonhuman animal, said animal having a nonyeast xenogeneic DNA segment of at least 100 kb stably integrated into the genome of all cells of said animal, said method comprising:

5
DI
cont
10 combining under fusing conditions embryonic stem cells of an available ES cell line derived from said animal and yeast spheroplasts, said spheroplasts containing a yeast artificial chromosome (YAC) comprising said xenogeneic DNA segment and including a marker for selection, whereby said xenogeneic DNA segment becomes integrated into the genome of said embryonic stem cells;

selecting for embryonic stem cells carrying said xenogeneic DNA segment by means of the marker;

15 transferring said embryonic cells into a host blastocyst and implanting said blastocyst in a pseudopregnant animal recipient, and allowing said blastocyst to develop to term to produce a chimeric animal carrying said xenogeneic DNA segment in the genome of at least some of the germ cells of
20 said animal; and

mating said chimeric animal with an animal of the same species to produce said modified animal carrying said xenogeneic DNA segment in all cells of said animal.

25
D2
5. (Amended) A method according to any of claims 1, 2 or 3, wherein said xenogeneic DNA is human DNA[, and/or wherein said xenogeneic DNA is immunoglobulin DNA in substantially intact form].

30
D3
6. (Amended) Embryonic stem cells comprising a genome modified according to the method of claims 1, 4, [or] 5 or 12.